

7. HYPERTENSION AND CORONARY HEART DISEASE

Dr Mirjana Cubrilo-Turek, MD, Ph.D.

Department of Internal Medicine, Sveti Duh General Hospital, Zagreb

Hypertension is the most prevalent treatable cardiovascular disease affecting approximately one in four adults or 140 million USA residents. It affects men and women in all socioeconomic groups equally. If untreated, hypertension is a major cause of stroke, coronary heart disease and renal failure as well as other conditions. Easily diagnosed, and in most instances readily controlled, hypertension is often unsuspected or inadequately treated.

The causes of high blood pressure are a bit of a mystery. According to the National Heart, Lung and Blood Institute, as many as 25% of adult Americans suffer from high blood pressure. Results of the Croatian national survey performed in 1997 in a sample of 5840 persons of both sexes, aged 18-65, showed about 28% adults to be hypertensive (BP>140/90 mm Hg), with a significantly greater prevalence recorded in men (32%) than in women (24%).

Major risk factors
Smoking
Dyslipidemia
Diabetes mellitus
Age >60 y
Sex (men and postmenopausal women)
Family history of cardiovascular disease: women <65 y or men <55 y
Target organ damage/Clinical cardiovascular disease
Heart diseases
Left ventricular hypertrophy
Angina or prior myocardial infarction
Prior coronary revascularization
Heart failure
Stroke or transient ischemic attack
Nephropathy
Peripheral arterial disease
Retinopathy

Table 1. Components of coronary risk stratification in patients with hypertension

7.1 The causes of high blood pressure

In most people, no specific cause of high blood pressure can be identified. It appears to be a distinct entity, due in part to a genetic predisposition (**essential hypertension**). **As much as 95% of the people with high blood pressure have essential hypertension. The probability of developing hypertension increases with age. More than half of all persons aged 65 are hypertensive. Treating high blood pressure significantly decreases the risk of stroke, myocardial infarction and cardiovascular death. Another type of high blood pressure called secondary hypertension is more common in younger individuals, usually due to some other problem (for example, persons taking birth control pills, renovascular patients, etc.).**

Category	Systolic	Diastolic
Optimal	<120	<80
Normal	<130	<85
High-normal	130-139	85-89
Grade 1 hypertension (mild)	140-159	90-99
Subgroup: borderline	140-149	90-94
Grade 2 hypertension (moderate)	160-179	100-109
Grade 3 hypertension (severe)	≥180	≥110
Isolated systolic hypertension	≥140	<90
Subgroup: borderline	140-149	

Table 2. Definition of hypertension according to the World Health Organization and the International Society of Hypertension (WHO/ISH) 1999

About 5% of patients requiring hypertension treatment can trace their high blood pressure to a physical cause such as kidney disease, however for 95% of patients who undergo hypertension treatment the causes of high blood pressure are unknown.

High risk groups. African Americans have a greater risk of developing hypertension than Caucasian Americans. Men are more likely to develop high blood pressure than women, and to develop hypertension at an earlier age. After the menopause, women become more susceptible to high blood pressure.

Category	Systolic (mm Hg)	Diastolic (mm Hg)
Optimal	<120 and	<80
Normal	<130 and	<85
High-normal	130-139 or	<85-89
Hypertension		
Stage 1	140-159 or	90-99
Stage 2	160-179 or	100-109
Stage 3	?180 or	?110

Table 3. Classification of blood pressure for adults aged ³ 18 according to Joint National Committee (JNC VI)

Hypertension is a major contributor to coronary heart disease and stroke, the two leading causes of death among adults worldwide. It is a silent epidemic affecting both sexes and rising with age. Hypertension, which affects approximately one in four adults in the USA, has an important role as an aetiological factor in the most prevalent treatable conditions such as cardiovascular disease. Untreated hypertension is a major cause of stroke, coronary heart disease (myocardial infarction), congestive heart failure and renal failure, as well as other conditions. Hypertension as a potentially serious condition produces various symptoms like sweating, palpitation, headache, and dizziness. Numerous scientific studies have estimated that between 55 and 70 million of the 140 million adults suffering from hypertension are unaware of it, and of those who know, fewer than half may be appropriately treated. Only 27% of all hypertensive patients in the USA are being treated appropriately. It is well known that the treatment of high blood pressure significantly decreases the risk of developing potentially fatal conditions.

In general, high blood pressure, or hypertension is defined as a systolic pressure of 140 mm Hg or greater and/or diastolic pressure of 90 mm Hg or greater, considered to be above the normal range. Definition of hypertension according to the guidelines of World Health Organization - International Society of Hypertension (1999) is presented in Table 2 and classification of blood pressure for adults according to Joint National Committee in Table 3.

7.2 Diagnosis

Hypertension is often referred to as a silent killer because the symptoms are often unnoticeable until complications occur. High blood pressure has a long asymptomatic period despite existing cardiovascular damage or kidney failure. The diagnosis of hypertension depends on careful blood pressure measurements and evaluation of the major risk factors. Two readings have to be taken together and averaged. Laboratory testing includes blood tests (haemoglobin and packed red cell volume, electrolytes, fasting blood glucose, serum lipids, serum creatinine levels, and serum uric acid), urine analyses (glycosuria, albuminuria), as well as evidence of the heart muscle thickening seen on electrocardiogram (cardiac rhythm, signs of

myocardial ischemia or myocardial infarction) and left ventricular hypertrophy or echocardiogram. Thorough history, physical examination and baseline laboratory data should be obtained in the patient whose blood pressure is usually above 140/90 mm Hg to assess the overall cardiovascular risk and to exclude identifiable causes of hypertension.

7.3 Risk stratification

In addition to blood pressure, JNC-VI recognizes the need to take into consideration other cardiovascular risk factors, target organ damage and coexistent cardiovascular disease to determine the overall risk profile of each patient and the need for drug therapy. In patients who have a low overall cardiovascular risk, even with a blood pressure as high as 160/100 mm Hg, lifestyle modifications alone are recommended initially. On the other hand, in patients at a high overall risk, such as those with diabetes, antihypertensive drug therapy is recommended even when the blood pressure is below 140/90 mm Hg. The treatment goal in these patients is a blood pressure of less than 130/85 mm Hg. In patients at risk, immediate drug therapy may be necessary.

7.4 Hypertension treatments

Today, there is a general agreement that not only severe but also mild-moderate hypertension should be treated. There is no full consensus on when to start antihypertensive treatment, and what is the first line drug treatment. The goal of antihypertensive therapy has to be individual in order to optimize the benefits and minimize the risk for each patient.

Antihypertensive drugs should be given initially in a relatively low dosage to avoid blood pressure lowering. The need to "start low and go slow" is even greater in the elderly, whose blood pressure should be reduced by gradual titration


The list of the more common approaches:

- Antihypertensive drugs
- Exercise
- Quitting smoking
- Regular blood pressure monitoring
- Stress reduction
- Dietary changes
- Weight loss
- Controlling cholesterol levels
- Reducing alcohol consumption

Lifestyle modifications for hypertension prevention and management:

- Lose weight if overweight

THRESHOLDS FOR INTERVENTION



* Unless malignant phase or hypertensive emergency, confirm over 1-2 weeks then treat.

** If cardiovascular complications, target organ damage, or diabetes is present, confirm over 3-4 weeks then treat; if absent re-measure weekly and treat if blood pressure persists at these levels over 4-12 weeks.

*** If cardiovascular complications, target organ damage, or diabetes is present, confirm over 12 weeks then treat; if absent re-measure monthly and treat if these levels are maintained and if estimated 10 year CHD risk is $\geq 15\%$

† Assessed with Cardiac Risk Assessor computer programme or coronary heart disease risk chart (6).

Figure 1. Algorithm for intervention according to blood pressure findings

- Limit alcohol intake to no more than 30 ml of ethanol (720 ml of beer, 300 ml of wine, or 60 ml of 100-proof whiskey) *per day* or 15 ml of ethanol *per day*
- Increase aerobic physical activity (30-45 min most days of the week)
- Reduce sodium intake to no more than 100 mmol/d (2.4 g of sodium or 6 g of sodium chloride)
- Maintain adequate intake of dietary potassium (approximately 90 mmol/d)
- Maintain adequate intake of dietary calcium and magnesium for general health
- Stop smoking and reduce intake of dietary saturated fat and cholesterol for overall cardiovascular health
- Diuretics
- Beta-blockers
- ACE inhibitors
- Calcium antagonists
- Alfa-blockers
- AT_1 -receptor blockers

Antihypertensive treatment now include six classes of antihypertensive agents according to WHO-ISH guidelines (1999):

Strong clinical evidence supports the “compelling indications,” as designated in the JNC-VI report, of certain drugs in four conditions that may accompany hypertension: diabetes with nephropathy (angiotensin converting enzyme [ACE] inhibitors); congestive heart failure (ACE inhibitors and diuretics); systolic hypertension in the elderly (diuretics and long-acting dihydropyridine calcium antagonists) and postmyocardial infarction (beta blockers with nonintrinsic sympathomimetic activity and with systolic dysfunction, ACE inhibitors).

Class of drug	Indication		Contraindications	
	Compelling	Possible	Possible	Compelling
a-blockers	Prostatism	Dyslipidemia	Postural hypotension	Urinary incontinence
ACE inhibitors	Heart failure Left ventricular dysfunction Type I diabetic nephropathy	Chronic renal disease* Type II diabetic nephropathy	Renal impairment* Peripheral vascular disease†	Pregnancy Renovascular disease
Angiotensin II receptor antagonists	Cough induced by ACE inhibitor‡	Heart failure Intolerance of other antihypertensive drugs	Peripheral vascular disease†	Pregnancy Renovascular disease
β-blockers	Myocardial infarction Angina	Heart failure§	Heart failure§ Dyslipidemia Peripheral vascular disease	Asthma or chronic obstructive pulmonary disease Heart block
Calcium antagonists (dihydropyridine)	Isolated systolic hypertension in elderly patients	Angina Elderly patients	-	-
Calcium antagonists (rate limiting)	Angina	Myocardial infarction	Combination with β blockade	Heart block Heart failure
Thiazides	Elderly patients	-	Dyslipidemia	Gout

Table 4. Compelling and possible indications and contraindications for the major classes of antihypertensive drugs (British Hypertension Society – Guidelines for Hypertension Management 1999)

*Angiotensin converting enzyme (ACE) inhibitors may be beneficial in chronic renal failure but should be used with caution. Close supervision and specialist advice are needed when there is established and significant renal impairment.

† Caution with ACE inhibitors and angiotensin II receptor antagonists in peripheral vascular disease because of association with renovascular disease.

‡ If ACE inhibitor indicated.

§ â-blockers may worsen heart failure, but in specialist hands may be used to treat heart failure.

HIGH RISK of death or nonfatal AMI. At least one of the following	INTERMEDIATE RISK of death or nonfatal AMI. No high-risk features plus one of the following	LOW RISK of death or nonfatal AMI. No high or intermediate features plus one of the following:
<ul style="list-style-type: none"> • Prolonged continuing pain not relieved by rest • Pulmonary edema • S₃ or rales • Hypotension with angina • Dynamic ST changes >1mm 	<ul style="list-style-type: none"> • Prolonged angina but resolved at time of evaluation (“stuttering”) • Rest angina lasting >20 min or relieved with nitroglycerin • Age >65 yrs • Dynamic T-wave changes • Q-waves with ST deviation ≤1 mm 	<ul style="list-style-type: none"> • Angina - increased in frequency, severity, or duration • Lower activity threshold before angina • New-onset angina >2 wk to 2 mo prior • Normal or unchanged ECG

(Modified from AHCPR. Clinical Practice Guideline - Unstable Angina.)

Table 5. Patients presenting with chest pain suggestive of ischemia: short-term risk of death. Based on clinical features and presenting electrocardiogram

The combinations of different antihypertensive drugs should be added until the goal blood pressure below 140/90 mmHg is reached, and even lower in those patients with renal insufficiency, diabetes or congestive heart failure

7.5 Coronary heart disease

Some 7 million Americans suffer from coronary heart disease (CHD), the most common form of heart disease. CHD is the number one killer of both men and women in the U.S. Each year, more than 500,000 Americans die of heart attacks caused by CHD.

Many of these deaths could be prevented because CHD is related to certain aspects of lifestyle. Risk factors for CHD include hypertension, high blood cholesterol, smoking, obesity, and physical inactivity - all of which can be controlled. Although medical treatments for heart disease have come a long way, the control of risk factors remains the key to preventing illness and death from CHD.

Risk factors that increase the risk of developing coronary heart disease:

Controllable:

- Hypertension
- High blood cholesterol
- Smoking
- Obesity
- Physical inactivity
- Diabetes
- Stress

Uncontrollable:

- Gender
- Heredity (family history of CHD)
- Age

Approximately 4 to 5 million patients will be evaluated in US emergency departments for chest pain. Of these patients 2 million will be diagnosed as having an acute ischemic syndrome. More than half a million of these patients will be hospitalized with a diagnosis of unstable angina, and 1.5 million will experience an acute myocardial infarction (AMI). Of the 1.5 million AMI patients, approximately half a million will die, and 50% of these deaths will occur within the first hour. Including prehospital mortality, the first prolonged attack of ischemic pain has a 34% fatality rate.

1.5.1 Symptoms of coronary heart disease

The term “acute coronary syndrome” has become commonly used to refer to patients presenting with ischemic chest pain. Chest pain (angina) or shortness of breath may be the earliest sign of CHD. The other clinical manifestations of CHD are: heaviness, tightness, pain, burning, pressure, or squeezing, usually behind the breastbone but sometimes also in the arms, neck, or jaws. Nevertheless, some people have heart attacks without ever having any of these symptoms.

Nomenclature of acute coronary syndromes (ACS):

- Unstable angina
- Non-Q-wave myocardial infarction
- Q-wave myocardial infarction

Initial assessment	Specific treatments
1 Targeted history, including AMI inclusions, thrombolytic exclusion 2 Vital signs and focused physical examination 3 12-lead ECG; serial ECGs as needed 4 Chest x-ray (preferably upright) 5 ECG monitoring	Reperfusion therapy: goals 10 Thrombolytic agents: door-to-needle time <30 min 11 Primary PTCA: door-to-dilation time <60 min Conjunctive therapy (combined with thrombolytic agents) 12 Aspirin 13 Heparin (especially with TPA) Adjunctive therapies 14 β -Adrenoceptor blockage if eligible 15 IV nitroglycerin (for anti-ischemic and antihypertensive effects) 16 ACE inhibitor (especially large anterior wall MI, heart failure without hypotension [SBP >100 mm Hg], previous MI)
Initial general treatment (memory aid: MONA)	
6 Morphine 2-4 mg q 5-10 min to provide adequate analgesia 7 Oxygen at 4 L/min; continue if arterial oxygen saturation <90% 8 Nitroglycerin sublingual or IV; tests for Prinzmetal's angina, reversible spasm; has anti-ischemic, antihypertensive effects 9 Aspirin 160-325 mg (chew and swallow)	

SBP indicates systolic blood pressure.

Table 6. Assessment and treatments of acute coronary syndrome

Discriminating among the acute coronary syndromes requires integration of clinical information, the amount and timing of cellular necrosis, examination of the cardiac markers (troponin I and T), and serial records of the 12-lead electrocardiogram (ECG).

(Modified from AHCPR. Clinical Practice Guideline - Unstable Angina.)

All of the coronary syndromes are initiated by the same event - rupture of an unstable, lipid-rich atheromatous plaque in an epicardial artery. Plaque disruption and fissuring activates platelet adhesion, fibrin clot formation, and coronary thrombosis. Without flow restriction these plaques do not cause clinical angina. Such plaques, however, have a lipid-rich core that makes them soft and prone to rupture. Inflammatory cells have been found in areas of plaque rupture. Stable plaques have less lipid and a thick cap that makes them more resistant to fissuring and formation of thrombi.

1.5.2 Diagnosis and tests for coronary heart disease

There is no single simple test for the diagnosis of CHD. Various diagnostic procedures have been used to establish CHD, after taking careful medical history and physical examination to determine its extent and severity, and to rule out other possible causes of symptoms.

An examination for CHD may include the following tests:

- New blood tests, which measure the level of particular enzyme markers (CK isoforms, troponin T and I, myoglobin) that rise with heart tissue damage, also help quickly diagnose a heart attack.

- Electrocardiogram (ECG) is a graphic record of the electrical activity of the heart. Abnormal heartbeats and some areas of damage, inadequate blood flow, and heart enlargement can be detected on the records.
- Exercise stress test (also called a treadmill test or exercise ECG) is used to record the heartbeat during exercise and blood pressure. In the test, an ECG is done before, during, and after exercising on a treadmill. ECG identifies any exercise stress-related abnormalities. Breathing rate may be measured as well. Exercise tests are useful but are not completely reliable; false positives (showing a problem where none exists) and false negatives (showing no problem when something is wrong) are fairly common.
- Nuclear scanning is sometimes used to show damaged areas of the heart and expose problems with the heart's pumping action. A small amount of radioactive material is injected into a vein, usually in the arm. A scanning camera records the nuclear material that is taken up by heart muscle (healthy areas) or not taken up (damaged areas).
- Coronary angiography (or arteriography) is a test used to explore coronary arteries by a series of x-rays. During an angiogram a fine, long flexible tube (catheter) is inserted into the artery of an arm or leg and passed through the tube into the arteries of the heart. The heart and blood vessels are then filmed while the heart pumps. Angiogram shows the flow within the coronary arteries and identifies any areas of narrowing or blockage caused by atherosclerosis.

7.6 Concepts of treatment and assessment

One of the major insights from the reperfusion era is that significant myocardial salvage is achieved not just from thrombolytic agents. Important benefits in long-term mortality and morbidity emerge from proper and early use

of oxygen, β -blockers, nitroglycerin, aspirin, heparin, and angiotensin-converting enzyme (ACE) inhibitors. From these insights has come the concept of three types of therapy:

- Reperfusion therapy - either thrombolytics or primary PTCA
- Adjunctive therapy - agents given instead of or in addition to thrombolytics
- Conjunctive therapy - agents given to enhance the benefits of thrombolytics

Modifying lifestyles in populations can have major protective effects against high blood pressure and various cardiovascular diseases. Lowering high blood pressure decreases death from stroke, coronary events and heart failure, slows progression of renal failure, prevents progression to severe complications of hypertension, and reduces all-cause mortality.

The goal of various international actions is to reduce morbidity, disability and premature mortality due to cardiovascular diseases with special attention to the prevention and control of hypertension by sharing knowledge, experience and technology in hypertension prevention among interested parties. Although medical treatments for heart disease have come a long way, the control of risk factors remains the key to preventing illness and death from coronary heart disease.

Recommended literature:

- 1 The Sixth Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC VI). Arch Intern Med 1997;157:2413-46. Copyright 1997, American Medical Association.
- 2 1999 World Health Organization - International Society of Hypertension.
- 3 Guidelines for the Management of Hypertension. J Hypertens 1999;17:151-83.
- 4 Blood Pressure 1999; 8 (Suppl 1):1-35
- 5 Turek S, Rudan I, Smolej-Narancic N, Szivovicza L, Cubrilo-Turek M, Žerjavic-Hrabak V, Rak-Kaic A, Vrhovski-Hebrang D, Prebeg Ž, Ljubicic M, Jancijevic B, Rudan P. A large cross-sectional study of health attitudes, knowledge, behaviour and risks in the post-war Croatian population. The First Croatian Health Project. Coll Antropol 2001; 25:77-96.
- 6 <http://bmj.com/cgi/content/full/319/7210/630/T2>
- 7 The Joint British Societies' Cardiac Risk Assessor computer programme and copies of the Joint British Societies' coronary heart disease risk assessment chart can be downloaded from the British Hypertension Society website (www.hyp.ac.uk/bhs).
- 8 Oparil S. Arterial hypertension. In: Cecil Textbook of Medicine 21st ed. Goldman L, Bennett JC. Philadelphia: W.B. Saunders Co., 2000; 258-73.